



Contents lists available at ScienceDirect

Bioresource Technology

journal homepage: www.elsevier.com/locate/biortech



A novel and robust recombinant *Pichia pastoris* yeast whole cell biocatalyst with intracellular overexpression of a *Thermomyces lanuginosus* lipase: Preparation, characterization and application in biodiesel production



Jinyong Yan^{*}, Xianliang Zheng, Shengying Li^{*}

Key Laboratory of Biofuels and Shandong Provincial Key Laboratory of Energy Genetics, Qingdao Institute of Bioenergy and Bioprocess Technology, Chinese Academy of Sciences, 189 Songling Road, Qingdao, Shandong 266101, China

HIGHLIGHTS

- A novel and robust *Pichia pastoris* yeast whole cell biocatalyst is reported.
- The short chain alcohol tolerant and easily scalable biocatalyst is developed.
- The biocatalyst gives improved biodiesel yield and productivity than Lipozyme TLIM.

ARTICLE INFO

Article history:

Received 2 September 2013
Received in revised form 8 October 2013
Accepted 11 October 2013
Available online 18 October 2013

Keywords:

Lipase
Whole cell catalyst
Pichia pastoris
Biodiesel
Waste cooking oils

ABSTRACT

A novel and robust recombinant *Pichia pastoris* yeast whole cell catalyst (WCC) with functional intracellular expression of *Thermomyces lanuginosus* lipase (Til) was constructed and characterized for biodiesel production from waste cooking oils. This permeabilized WCC was able to convert waste cooking oils to biodiesel with 82% yield within 84 h at 6% dosage whole cells. The WCC showed two fold catalytic activity of 0.73 U/mg DCW compared to its commercial counterpart Lipozyme TLIM (immobilized Til). Short chain alcohol tolerance of this WCC was significantly improved compared to Lipozyme TLIM. This beneficial property enabled it to catalyze biodiesel production efficiently with one step addition of methanol. The reusability of this biocatalyst retained 78% activity after three batch cycles. This easily prepared and cost-effective WCC showed better catalytic performance than Lipozyme TLIM with respect to biodiesel yield and productivity, thus suggesting a promising cost-effective biocatalyst for biodiesel production.

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

The fatty acid methyl esters (FAMES) and fatty acid ethyl esters (FAEEs) produced by transesterification of triglycerides from vegetable oils and animal fats, known as biodiesel, have been receiving ever-increasing attentions because they are renewable and environment-friendly biofuels with excellent fuel properties (Tan et al., 2010). The cost of biodiesel production is mainly dependent on the price of oil feedstocks. The high price of edible oils makes their conversion into biodiesel economically infeasible (Yan et al., 2011). Instead, the low cost waste cooking oils are a promising

alternative to edible oils for biodiesel production despite of their low quality.

For common biodiesel production using refined vegetable oils as feedstock, alkaline catalysis has been proven much more efficient and cost-effective than acid catalysis. However, the high content free fatty acids (FFAs) present in waste cooking oils need to be reduced by acid-catalyzed esterification prior to base-catalyzed transesterification. Otherwise, the soap formation between FFAs and bases would deactivate the alkali catalysts to hamper the transesterification (Lam et al., 2010). In addition, the use of both acids and bases could lead to a number of significant problems such as reactor corrosion, complex neutralization, and environmental pollution (Lam et al., 2010). By comparison, some lipases are able to efficiently catalyze esterification of FFAs and transesterification of triglycerides simultaneously in presence of methanol or ethanol (Yan et al., 2011; Adachi et al., 2013a,b), thus representing suitable biocatalysts for generation of biodiesel from waste

Abbreviations: WCCs, whole cell catalysts; FAMES, fatty acid methyl esters; FAEEs, fatty acid ethyl esters; FFAs, free fatty acids; DCW, dry cell weight; BSPs, biomass support particles.

^{*} Corresponding authors. Tel.: +86 532 80662675; fax: +86 532 80662778 (J. Yan).

E-mail addresses: yjiny@126.com (J. Yan), lishengying@qibebt.ac.cn (S. Li).

cooking oils. Moreover, other advantages related to enzymatic biodiesel production include mild reaction conditions, less environment impacts and easy separation of byproduct glycerol (Yan et al., 2011).

Apart from these advantageous catalytic properties of lipases, the cost of lipases is another important factor to be taken into account with respect to economic biodiesel production. To date, three major types of lipase biocatalysts, including free enzymes, immobilized enzymes and whole cell catalysts (WCCs) have been employed for synthesis of biodiesel. Free enzymes are unstable, difficult to be reused and hence expensive. Immobilized lipase preparations have been widely used in many industrial processes including biodiesel production. Various immobilization carriers and immobilization technologies have offered opportunities in tuning enzyme properties such as stability, activity and selectivity, as well as high volumetric loading (Barbosa et al., 2013; Rodrigues et al., 2013; Verma et al., 2013). However, they are expensive since their preparation requires costly steps of enzyme extraction and immobilization (Tan et al., 2010). To achieve an applicable immobilized lipase product, a lot of work for seeking optimal immobilization carriers and immobilization conditions is necessary. These complicate the use of immobilized lipases for industrial production of biodiesel. In comparison, the lipase WCC may provide the best solution to reduce the biocatalyst cost because its simple preparation does not require any purification and immobilization steps (Fukuda et al., 2008). Similar to immobilized enzymes, the stability and reusability of lipases in the form of WCC can be significantly improved compared to free enzymes (Takaya et al., 2011; Koda et al., 2010). Taken together, the WCC lipase represents a promising biocatalyst for biodiesel production in a cost-effective manner.

So far, three major forms of WCCs have been applied in biodiesel production. First, the fungal cells of *Rhizopus* spp with cell-bound lipases represent examples of natural WCCs (Li et al., 2007; He et al., 2008). Specifically, *Rhizopus oryzae* and *Rhizopus chinensis* were found to secrete a portion of lipases into extracellular space, while retain the rest as the intracellular membrane-bound form. Apparently, not all lipases exist as WCC in this case, and additional immobilization with biomass support particles (BSPs) or glutaraldehyde is often preferred for improvement of their stability and activity (Ban et al., 2002; Xiao et al., 2011).

Second, display of lipases on the surface of yeast cells including *Pichia pastoris* and *Saccharomyces cerevisiae* via expression of genetically fused enzyme-anchor proteins has been developed as another type of WCC for biodiesel production (Matsumoto et al., 2002; Huang et al., 2012). These yeast cells with surface displayed lipases are more substrate accessible compared to intracellular enzymes. However, the expression level and enzymatic activity of the anchored lipases are generally low. In addition, establishment of the compatible relationship among anchor protein, target lipase and host cell is not easy work.

Third, the strategy of preparing lipase WCCs for biodiesel production by intracellular overexpression of lipases inside heterologous host has emerged recently. The investigations were carried out in *Escherichia coli* (Yan et al., 2012; Li et al., 2012; Gao et al., 2009) and *S. cerevisiae* (Matsumoto et al., 2001). Surprisingly, this strategy has not yet been extended to the yeast of *P. pastoris*, which can achieve higher cell density, and a greater expression level of functional heterologous enzymes due to less hyper-glycosylation than *S. cerevisiae* yeast (Patrick et al., 2005). It is able to accumulate large amounts of intracellular enzymes to the level as high as that of excreted proteins (Patrick et al., 2005), thereby becoming a “self-immobilized” biocatalyst that retains almost all active enzymes inside the cells.

Thermomyces lanuginosus lipase (Tll) has been well known as a versatile biocatalyst in various industrial areas including biodiesel production, modification of oils, chiral resolution of racemic

compounds and others (Fernandez-Lafuente, 2010). Thus, in the present work, we selected this useful enzyme to exploit its new preparation as the form of WCC, and applied it into biodiesel production. Specifically, a recombinant *P. pastoris* yeast strain with intracellular expression of *T. lanuginosus* lipase was constructed for preparation of a novel WCC. The organic solvent tolerance and thermostability of this WCC were characterized. Its ability of converting waste cooking oils into biodiesel was studied. To the best of our knowledge, this is the first example of using recombinant *P. pastoris* yeast to functionally overexpress intracellular lipases as a robust WCC for biodiesel production.

2. Methods

2.1. Plasmid, strains, media and reagents

The plasmid pPICZA and yeast strain *P. pastoris* X-33 were purchased from Invitrogen. *E. coli* DH5 α was reserved in our laboratory. Pretreated waste cooking oils (a mixture of waste oils from kitchen sewer) with 19% FFA content were kindly donated by Lyming Environmental Protection Technology Co., Ltd. (Shanghai, China), and their main properties were as follows: saponification value of 366 mg KOH/g, acid value of 170 mg KOH/g, density of 0.89 g/cm³, water content of 0.1%, and mechanical impurities of 0.4%. Immobilized enzyme Lipozyme TLIM was the product of Novozymes. Standard FAME mix (equal mole of methyl palmitate, methyl stearate, methyl oleate, methyl linoleate, and methyl linolenate) and internal standard (*n*-hexadecane) were purchased from Sigma–Aldrich. All used restricted enzymes and genetic manipulation kits were supplied by Takara. The antibiotic zeocin was bought from Solarbio Science & Technology Co., Ltd (Beijing, China). Yeast nitrogen base (YNB), yeast extract, tryptone and biotin were from Oxoid. Organic solvents and other reagents were analytical grade.

Recipes for used media were as follows: YPDS medium: 2% tryptone, 1% yeast extract, 1% dextrose, 1 M sorbitol, 1.5% agar, supplemented with certain concentration of zeocin. BMGY medium: 2% tryptone, 1% yeast extract, 100 mM potassium phosphate buffer, pH 7.0, 1.34% YNB, 0.4 ppm biotin, 1% (v/v) glycerol. BMMY medium: 2% tryptone, 1% yeast extract, 100 mM potassium phosphate buffer, pH 7.0, 1.34% YNB, 0.4 ppm biotin, 1% (v/v) methanol. Low salt LB medium: 1% tryptone, 0.5% yeast extract, 0.5% NaCl, supplemented with 25 μ g/ml zeocin if required, 1.5% agar powder was added for agar plate preparation.

2.2. Construction of plasmid and recombinant strain

The codon optimized *tll* gene (Genbank Accession No. AF054513.1) based on the codon bias of *P. pastoris* (Supporting information, Fig. S1) was synthesized by GenScript. Using the synthetic gene as template, the *tll* fragment with its own signal peptide removed and an ATG start codon added before the mature lipase sequence was amplified by PCR with primers as follow: forward, 5'-CCGGAATTCATGTCACCTATCAGAAGAGAAGTTTCAC-3' (the underlined nucleotides indicate the *EcoRI* restriction site); reverse, 5'-AAGGAAAAAAGCGGCGCCTACAAGCAAGTTCCGATAAGACC-3' (the underlined bases denote the *NotI* cutting site). Next, the *EcoRI* and *NotI* double digested *tll* fragment was ligated into the *EcoRI*/*NotI* pretreated plasmid pPICZA. The resulted plasmid was transformed into *E. coli* DH5 α competent cells. Transformants grown on low salt LB medium containing 25 μ g/ml zeocin was employed for preparation of pPICZA-*tll*, whose sequence was confirmed by DNA sequencing at Sangon, Shanghai.

For yeast transformation, pPICZA-*tll* was linearized by *SacI* and transformed into competent cells of *P. pastoris* X-33 via electroporation under the following conditions: 0.2 cm cuvette, 1500 V,

200 Ω , 25 μ F. The positive transformants harboring chromosomal integration of TII expression cassettes were preliminarily selected on YPDS plates containing 100 μ g/ml zeocin. Multi-copy integrated recombinants were further screened on YPDS plates supplemented with gradually increased concentration of zeocin (500–2000 μ g/ml).

2.3. Intracellular expression of recombinant TII lipase

A single colony of *P. pastoris* containing multi-copy TII expression cassettes was inoculated into 5 ml BMGY medium and cultured at 28 °C, 250 rpm, for 24 h. This seed culture was transferred to 50 ml BMGY medium in a 500 ml baffled flask and grown until OD₆₀₀ reaching 4–6. The cells were sterilely collected by centrifugation and re-suspended in a 2 l baffled flask by 400 ml pre-loaded fresh BMMY medium. Subsequently, 1% (v/v) methanol was added into culture every 24 h for induction of lipase expression. The strain harboring empty vector pPICZA was subject to the same manipulation as control for expression analysis.

2.4. Determination of cell density and catalytic activity

At designated time intervals, 0.5 ml aliquots of induced cell culture in BMMY medium were withdrawn for determination of cell density by measuring OD₆₀₀. Yeast cells with intracellular expression of TII were collected from 5 ml culture by centrifugation, re-suspended and washed twice with 0.85% (w/v) NaCl solution, re-centrifuged, and used as untreated whole cell biocatalysts. Following the previously established protocol (Yan et al., 2007) with slight modifications, the hydrolytic reactions were conducted using emulsified olive oil as substrate. Emulsified olive oil was prepared by homogenizing a mixture of olive oil and 2% polyvinyl alcohol solution (1:3, v/v) using homogenizer at 6000 rpm for 10 min. The reaction mixture contained 5 ml of 50 mM Tris–HCl buffer (pH 8.0), 4 ml of emulsified olive oil as substrate, and 50 mg whole cells or Lipozyme TLIM as catalysts. The hydrolysis reactions were carried out at 40 °C and pH 8.0 for 10 min, and stopped by addition of 15 ml ethanol. Activity was determined titrimetrically using 50 mM NaOH as titrant. The same amount of whole cell lipases was inactivated by boiling at 100 °C for 1 h and used as control. One unit of lipase activity was defined as the amount of enzyme that hydrolyzes olive oils to form 1 μ mol free fatty acid per minute under testing conditions. The specific activity was calculated as units of enzyme activity possessed by one milligram of dry cell weight (U/mg DCW). DCW was determined by oven drying wet cells to constant weight at 60 °C. With respect to the commercial immobilized enzyme Lipozyme TLIM, its activity was defined as units of enzyme activity held by one milligram of solid catalysts. Unless otherwise specified, all quantitative experiments in this study were performed in duplicate. $P < 0.05$ indicated statistical significance (*).

2.5. Characterization of organic solvent tolerance and thermostability

The untreated yeast whole cell biocatalysts were incubated with 25% (v/v) short chain alcohol (methanol, ethanol or isopropyl alcohol) at room temperature for 30 min or heated at 50–70 °C for 30 min. Organic solvent tolerance and thermostability were characterized by determining residual activity of whole cell catalysts after alcohol treatment and heating, respectively, relative to that of untreated cells. Similarly, these properties of Lipozyme TLIM were determined under the same conditions.

2.6. Permeabilization of yeast whole cells

Freeze drying permeabilization was conducted by freezing yeast cells at –80 °C for 12 h and drying with lyophilizer for 48 h. Isopropyl alcohol treatment coupled to freeze-drying permeabilization was performed by incubation of cells in 25% isopropyl alcohol for 30 min at 4 °C and followed by freeze-drying.

2.7. Biotransformation of waste cooking oils into biodiesel

Reaction mixture containing 2 g waste cooking oils and total 300 μ l methanol (molar ratio of 1:4) was emulsified with 5% (wt% relative to oil) polyvinyl alcohol solution and placed in a conical flask with stopper. 6% (wt% relative to oil) whole cell lipase or 12% (wt% relative to oil) immobilized lipase Lipozyme TLIM were added to initiate the biotransformation. One third of total amount of methanol was added at 0, 24 and 48 h (0, 12 and 24 h for Lipozyme TLIM). To investigate the methanol tolerance of biocatalysts, all methanol was added at once at 0 h to reaction mixture. Reactions were carried out at 40 °C, 250 rpm with a heating magnetic stirring system. The whole cells harboring empty vector were subject to the same preparation procedure and reaction conditions as control.

To evaluate the reusability of biocatalysts, the yeast whole cell catalysts were rescued by centrifugation at 6000 \times g for 5 min after every batch reaction. Upon washing with tert-butanol and drying, the recovered biocatalysts were used for new batch reaction under the same conditions. The operational stability in each batch cycle was characterized by measuring the relative biodiesel yield compared to that of the first reaction.

2.8. Identification of FAMES

The aliquots taken from reaction mixture at certain time points were mixed with 0.1 ml deionized water and centrifuged for 10 min at 11,000 \times g. Then, 10 μ l top layer was mixed with 990 μ l *n*-hexane containing 1 mM internal standard of *n*-hexadecane as the testing sample for GC analysis. Briefly, 1 μ l sample was injected into Agilent 7890A GC system equipped with flame-ionization detector and a capillary column (INNOWAX, Agilent, 30 m \times 0.25 mm \times 0.25 μ m) for FAME analysis. The column temperature was increased from 150 to 260 °C at 5 °C/min, and kept at 260 °C for 5 min. The temperatures of injector and detector were set at 240 and 280 °C, respectively. The produced FAMES were identified through comparison with corresponding authentic standards regarding their retention time. Quantitative analysis of FAMES was performed as described previously (Li et al., 2012) using internal standard *n*-hexadecane as reference.

3. Results and discussion

3.1. Recombinant strain construction and whole cell biocatalyst preparation

The recombinant *P. pastoris* yeast strain of X-33 (Mut⁺) with intracellular overexpression of TII was constructed through multiple chromosomal integration of the expression cassette consisting of the *tII* gene with signal peptide sequence removed and the selectable zeocin resistance gene (*zeo*^r) onto the AOX1 (alcohol oxidase 1) locus of host genome. This was achieved by multiple single crossover events occurred at the 5' AOX1 promoter region or the AOX1 transcription terminator region via homologous recombination between linearized vector and host genome (Supporting information, Fig. S2). Positive transformants harboring multiple integrated TII expression cassettes were stepwise

screened on YPDS plates supplemented with gradually increased concentration (500–2000 µg/ml) of zeocin.

A recombinant *P. pastoris* yeast strain containing multi-copy TII-*zeo^r* gene cassettes demonstrated by good resistance to 2000 µg/ml of zeocin was selected for lipase fermentation in shaking flask. The general procedure for expression of recombinant lipases in *Pichia* yeast consists of two stages: (1) the biomass accumulation in BMGY medium to avoid the potential toxicity resulted from heterogenous protein and/or the expression inducer methanol; (2) the lipase production after the cell density reached OD₆₀₀ of 4–6 in BMGY medium, at which the yeast cells were transferred from BMGY to BMMY medium for further biomass production and lipase expression. As shown in Fig. 1A, the peak cell density reached OD₆₀₀ of 52 at 48 h after methanol induction. The high cell density is a significant advantage with regard to cost-effective preparation of biocatalysts at large scale. More importantly, the highest specific olive oil hydrolyzing activity of the whole cell TII catalyst was determined to be 0.73 U/mg DCW, which is significantly higher than 0.37 U/mg solid catalyst of the commercially available immobilized TII-Lipozyme TLIM (Fig. 1B). This strongly suggests that the yeast whole cells are likely to be a better “immobilizing carrier” for functional TII enzymes than the poly acrylic resin used in Lipozyme TLIM. If further taking simpler and cheaper preparation into account, TII WCC is a promising candidate to be developed into an industrial biocatalyst.

3.2. Short chain alcohol tolerance of TII WCC

Intolerance of lipase enzymes toward short chain alcohols would limit their application in biodiesel production. To investigate whether TII WCC is tolerant to short chain alcohols, methanol, ethanol and isopropyl alcohol (25%, v/v) were used to pre-treat the yeast whole cells for 30 min, and the residual activity of hydrolyzing olive oils was measured. Interestingly, the alcohol treated WCCs displayed higher activity than the untreated cells (Fig. 2A). In particular, a 45% improvement of activity was observed in the case of isopropyl alcohol treatment. These results are consistent with the previous study that alcohol treatment of *S. cerevisiae* cells enhanced the conversion rate of *S*-lactoylglutathione (Kondo et al., 2000). On the contrary, the short chain alcohol treatments have adverse effects on the commercial immobilized enzyme Lipozyme TLIM, which are in good agreement with a previous report (Martín and Otero, 2008).

The comprehensive effects imposed by short chain alcohol treatment could be categorized into three positive or negative aspects including (1) the polar microenvironment created by short chain alcohol for formation of essential water layer to stabilize

the active enzymes (Rodrigues et al., 2013); (2) the role as cell permeabilizers, which facilitate substrates across yeast cell walls and cell membranes to improve substrate-enzyme interactions; and (3) the enzyme deactivation effect by short chain alcohol. It is particularly noteworthy that the intracellular environment of WCC is likely to be a significant factor for protecting inside lipase from deactivating alcohol and for maintaining active enzyme conformation. In contrast, the same alcohol treatment led to adverse effect on immobilized lipase Lipozyme TLIM, thus highlighting this significant advantage of WCC TII over Lipozyme TLIM with respect to short chain alcohol tolerance.

3.3. Thermal stability of TII WCC

TII has been known as a thermophilic enzyme (Fernandez-Lafuente, 2010). To evaluate whether the form of biocatalyst alters the thermostability of TII, the WCC activity to hydrolyze olive oils was measured at a range of high temperatures. Similar to Lipozyme TLIM, TII WCC exhibited high thermal stability (Fig. 2B), retaining 97% and 89% of activity after incubation at 50 and 60 °C for 30 min, respectively. Despite a significant loss of activity at 70 °C, it was still more stable than Lipozyme TLIM.

3.4. Biotransformation of waste cooking oils to biodiesel by TII WCC

Given that TII WCC demonstrated significantly higher short chain alcohol tolerance than the immobilized enzyme Lipozyme TLIM, we expected that it should be a better biocatalyst in short chain alcohols involved in biodiesel production. Thus, using Lipozyme TLIM as reference, the activity of TII WCC to transform waste cooking oils into FAMES was assessed. As shown in Fig. 1B, TII WCC showed the olive oil hydrolyzing activity of 0.73 U/mg DCW, which is almost two-fold that of Lipozyme TLIM (0.37 U/mg catalyst). Therefore, the amount of immobilized lipase (12% wt) was doubled relative to WCC (6% wt) in order to apply the same level of enzymatic activity into biotransformation of waste cooking oils to biodiesel.

Unexpectedly, the maximal biodiesel yield of TII WCC (44% at 108 h, Fig. 3) turned out to be approximately 60% of that of Lipozyme TLIM (75% at 48 h). The reason could be limited substrate accessibility to intracellular lipases due to the shielding effect of yeast cell walls and cell membranes. Thus, permeabilization of yeast cells could promote mass transfer across these physical barriers to improve the catalytic performance of TII WCC. To test this hypothesis, freeze-drying permeabilization was first attempted since it has been known as a simple but effective permeabilization method for improvement of biotransformation by *S. cerevisiae*




Fig. 1. (A) Time course of cell density and the specific olive oil hydrolyzing activity of TII WCC. (B) The specific olive oil hydrolyzing activity of TII WCC and its commercial counterpart Lipozyme TLIM (immobilized TII).




Fig. 2. Effects of short chain alcohols (A) and temperature (B) on TII WCC and Lipozyme TLIM.




Fig. 3. Biodiesel production from waste cooking oils by variant forms of TII.




Fig. 4. Biodiesel production from waste cooking oils by TII WCC and Lipozyme TLIM in the reaction with one step methanol addition.

yeast cells (Matsumoto et al., 2001). Using the freeze-dried TII WCC, the highest biodiesel yield was increased to 71% at 108 h. According to above described results (Fig. 2A), alcohol treatment is another effective method of permeabilization. Upon a combination of isopropyl alcohol (the best permeablizing alcohol among the tested short chain alcohols) treatment and freeze-drying, the maximal biodiesel yield of TII WCC was further enhanced to 82% at 84 h, which is higher than that of Lipozyme TLIM despite of longer reaction time.

It is evident that Lipozyme TLIM showed much higher initial catalytic rate than TII WCC, consistent with the previous result from Matsumoto et al. This could be explained by more efficient mass transfer in immobilized TII. After all, the permeabilization of WCC would take extra time. However, TII WCC demonstrated higher biodiesel yield and biodiesel productivity (9.76% yield/g catalyst·h) provided with additional reaction time of 36 h compared to Lipozyme TLIM (7.81% yield/g catalyst·h). Importantly, it is much easier to prepare large amounts of TII WCC by cultivation, cell harvesting, and permeabilization treatment.

Enzyme inactivation by methanol is one of central limitations in lipase mediated biodiesel production. Although stepwise addition of methanol has solved this problem to some extent (Li et al., 2007), one step addition of methanol is still preferred because it would simplify the process for practical application, especially at industrial scale. Thus, the compatibility of TII WCC with one step addition of methanol was evaluated. Compared to Lipozyme TLIM, the yeast cells permeabilized by isopropyl alcohol treatment and freeze-drying displayed significantly higher biodiesel yield in the reaction with one step methanol

addition (Fig. 4). This is likely due to its much better tolerance to methanol.

Taken together, TII WCC is probably a better option for economic production of biodiesel from waste cooking oils than costly immobilized Lipozymes. However, immobilized lipases may still hold advantages in other applications such as chiral resolution of racemic compounds and synthesis of high value-added fine chemicals. Aiming to develop it into an applicable biocatalyst, the systematic optimization of parameters including catalyst dosage, substrate molar ratio, water content, reaction temperature and reaction time involved in biodiesel production is currently ongoing in our laboratory.

3.5. Recyclable stability of TII WCC and immobilized Lipozyme TLIM

Biocatalysts with good recyclable stability further reduce the cost of biotransformation. The permeabilized TII WCC by isopropyl alcohol treatment and freeze-drying was used to catalyze biodiesel synthesis in continuous batches. As result, TII WCC retained more than 70% activity after three biotransformation cycles (Fig. 5), showing comparable reusability to Lipozyme TLIM. It is worth noting that no lipase activity was detected in the washing organic solvents, which indicates the yeast cell walls and/or cell membranes serve well as natural barriers to keep all biocatalysts inside cells. The similar activity loss of the two forms of TII in recyclable batches might be attributed to gradual deactivation by methanol and various negative components in waste cooking oils such as metal ions and glue substances.




Fig. 5. Biodiesel production by TII WCC and Lipozyme TLIM in batch cycles.

4. Conclusion

A novel and robust recombinant *P. pastoris* yeast whole cell biocatalyst with intracellular overexpression of TII was developed for biodiesel production from waste cooking oils. The permeabilization of TII WCC led to 82% biodiesel yield, and remained 78% activity after three batch cycles. It also efficiently catalyzed biodiesel production in the reaction with one step methanol addition. This catalytically efficient, thermostable, organic solvent tolerant, easily scalable, reusable, and cost-effective TII WCC holds great potential for economic biodiesel production. The yeast whole cell biocatalytic system also establishes a new platform for lipase-associated biocatalysis and biotransformation processes.

Acknowledgements

This work was partially supported by funding from the National High Technology Research and Development Program of China (NO. SS2014AA091610) and “Recruitment Program of Global Experts, 2012” (S. Li). We are grateful to Ms. Wei Zhou at Lvming Environmental Protection Technology Co., Ltd. (Shanghai, China) for providing us with the waste cooking oils.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.biortech.2013.10.037>.

References

Adachi, D., Koh, F.H., Hama, S., Ogino, C., Kondo, A., 2013a. A robust whole-cell biocatalyst that introduces a thermo- and solvent-tolerant lipase into *Aspergillus oryzae* cells: characterization and application to enzymatic biodiesel production. *Enzyme Microb. Technol.* 52, 331–335.

Adachi, D., Hama, S., Nakashima, K., Bogaki, T., Ogino, C., Kondo, A., 2013b. Production of biodiesel from plant oil hydrolysates using an *Aspergillus oryzae* whole-cell biocatalyst highly expressing *Candida antarctica* lipase B. *Bioresour. Technol.* 135, 410–416.

Ban, K., Hama, S., Nishizuka, K., Kaieda, M., Matsumoto, T., Kondo, A., Noda, H., Fukuda, H., 2002. Repeated use of whole-cell biocatalysts immobilized within biomass support particles for biodiesel fuel production. *J. Mol. Catal. B Enzym.* 17, 157–165.

Barbosa, O., Torres, R., Ortiz, C., Berenguer-Murcia, A., Rodrigues, R.C., Fernandez-Lafuente, R., 2013. Heterofunctional supports in enzyme immobilization: from traditional immobilization protocols to opportunities in tuning enzyme properties. *Biomacromolecules* 14, 2433–2462.

Fernandez-Lafuente, R., 2010. Lipase from *Thermomyces lanuginosus*: uses and prospects as an industrial biocatalyst. *J. Mol. Catal. B Enzym.* 62, 197–212.

Fukuda, H., Hama, S., Tamalampudi, S., Noda, H., 2008. Whole-cell biocatalysts for biodiesel fuel production. *Trends Biotechnol.* 26, 668–673.

Gao, B., Su, E.Z., Lin, J.P., Jiang, Z.B., Ma, Y.S., Wei, D.Z., 2009. Development of recombinant *Escherichia coli* whole-cell biocatalyst expressing a novel alkaline lipase-coding gene from *Proteus* sp. for biodiesel production. *J. Biotechnol.* 139, 169–175.

He, Q., Xu, Y., Teng, Y., Wang, D., 2008. Biodiesel production catalyzed by whole-cell lipase from *Rhizopus chinensis*. *Chin. J. Catal.* 29, 41–46.

Huang, D.F., Han, S.Y., Han, Z.L., Lin, Y., 2012. Biodiesel production catalyzed by *Rhizomucor miehei* lipase-displaying *Pichia pastoris* whole cells in an isooctane system. *Biochem. Eng. J.* 63, 10–14.

Kondo, A., Liu, Y., Furuta, M., Fujita, Y., Matsumoto, T., Fukuda, H., 2000. Preparation of high activity whole cell biocatalyst by permeabilization of recombinant flocculent yeast with alcohol. *Enzyme Microb. Technol.* 27, 806–811.

Koda, R., Numata, T., Hama, S., Tamalampudi, S., Nakashima, K., Tanaka, T., Ogino, C., Fukuda, H., Kondo, A., 2010. Ethanolysis of rapeseed oil to produce biodiesel fuel catalyzed by *Fusarium heterosporum* lipase-expressing fungus immobilized whole-cell biocatalysts. *J. Mol. Catal. B Enzyme* 66, 101–104.

Li, W., Du, W., Liu, D.H., 2007. *Rhizopus oryzae* IFO 4697 whole cell catalyzed methanolysis of crude and acidified rapeseed oils for biodiesel production in tert-butanol system. *Process Biochem.* 42, 1481–1485.

Li, A.T., Ngo, T.P.N., Yan, J.Y., Tian, K.Y., Li, Z., 2012. Whole-cell based solvent-free system for one-pot production of biodiesel from waste grease. *Bioresour. Technol.* 114, 725–729.

Lam, M.K., Lee, K.T., Mohamed, A.R., 2010. Homogeneous, heterogeneous and enzymatic catalysis for transesterification of high free fatty acid oil (waste cooking oil) to biodiesel: a review. *Biotechnol. Adv.* 28, 500–518.

Matsumoto, T., Fukuda, H., Ueda, M., Tanaka, A., Kondo, A., 2002. Construction of yeast strains with high cell surface lipase activity by using novel display systems based on the Flo1p flocculation functional domain. *Appl. Environ. Microbiol.* 68, 4517–4522.

Matsumoto, T., Takahashi, S., Kaieda, M., Ueda, M., Tanaka, A., Fukuda, H., Kondo, A., 2001. Yeast whole-cell biocatalyst constructed by intracellular overproduction of *Rhizopus oryzae* lipase is applicable to biodiesel fuel production. *Appl. Microbiol. Biotechnol.* 57, 515–520.

Martin, E.H., Otero, C., 2008. Different enzyme requirements for the synthesis of biodiesel: Novozym 435 and Lipozyme TLIM. *Bioresour. Technol.* 99, 277–286.

Patrick, S.M., Fazenda, M.L., McNeil, B., Harvey, L.M., 2005. Heterologous protein production using the *Pichia pastoris* expression system. *Yeast* 22, 249–270.

Rodrigues, R.C., Ortiz, C., Berenguer-Murcia, A., Torres, R., Fernández-Lafuente, R., 2013. Modifying enzyme activity and selectivity by immobilization. *Chem. Soc. Rev.* 42, 6290–6307.

Takaya, T., Koda, R., Adachi, D., Nakashima, K., Wada, J., Bogaki, T., Ogino, C., Kondo, A., 2011. Highly efficient biodiesel production by a whole-cell biocatalyst employing a system with high lipase expression in *Aspergillus oryzae*. *Appl. Microbiol. Biotechnol.* 90, 1171–1177.

Tan, T.W., Lu, J.K., Nie, K.L., Deng, L., Wang, F., 2010. Biodiesel production with immobilized lipase: a review. *Biotechnol. Adv.* 28, 628–634.

Verma, M.L., Barrow, C.J., Puri, M., 2013. Nanobiotechnology as a novel paradigm for enzyme immobilisation and stabilisation with potential applications in biodiesel production. *Appl. Microbiol. Biotechnol.* 97, 23–39.

Xiao, M., Qi, C., Obbard, J.P., 2011. Biodiesel production using *Aspergillus niger* as a whole cell biocatalyst in a packed-bed reactor. *GCB Bioenergy* 3, 293–298.

Yan, J.Y., Yang, J.K., Xu, L., Yan, Y.J., 2007. Gene cloning, overexpression and characterization of a novel organic solvent tolerant and thermostable lipase from *Galactomyces geotrichum* Y05. *J. Mol. Catal. B Enzym.* 49, 28–35.

Yan, J.Y., Yan, Y.J., Liu, S.X., Hu, J., Wang, G.L., 2011. Preparation of cross-linked lipase-coated micro-crystals for biodiesel production from waste cooking oil. *Bioresour. Technol.* 102, 4755–4758.

Yan, J.Y., Li, A.T., Xu, Y., Ngo, T.P.N., Phua, S.C., Li, Z., 2012. Efficient production of biodiesel from waste grease: one-pot esterification and transesterification with tandem lipases. *Bioresour. Technol.* 123, 332–337.